



HAL
open science

Quantifying perinatal brain maturation using anatomical MRI

Guillaume Auzias, François Rousseau, Sylvain Takerkart, Nadine Girard, Christine Deruelle, Olivier A Coulon, Julien Lefèvre

► **To cite this version:**

Guillaume Auzias, François Rousseau, Sylvain Takerkart, Nadine Girard, Christine Deruelle, et al.. Quantifying perinatal brain maturation using anatomical MRI. 2019 41st Annual International Conference of the IEEE Engineering in Medicine and Biology Society (EMBC), 2019, Berlin, Germany. hal-02286005

HAL Id: hal-02286005

<https://imt-atlantique.hal.science/hal-02286005v1>

Submitted on 13 Sep 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Quantifying perinatal brain maturation using anatomical MRI

G. Auzias, F. Rousseau, S. Takerkart, N. Girard, C. Deruelle, O. Coulon, J. Lefèvre

Abstract—Perinatal Magnetic Resonance Imaging has proven to be suitable and efficient for detecting alterations in a growing number of pathologies. Major advances in brain MRI acquisition and processing open the way to quantitative analyses not achievable so far. Recent results highlight the great potential of regional descriptors of the geometry of the cortical surface extracted from anatomical MRI to characterize typical and abnormal maturation trajectories in the perinatal period.

I. INTRODUCTION

Continuous, sustained methodological advances in image acquisitions and processing (fast acquisition, movement correction, super-resolution/reconstruction) allow to address new problems that were not affordable few years ago (e.g. RS-fMRI, tractography) [1], [2]. More specifically, dedicated image processing pipelines allow to extract a 3D mesh representation of the cortical surface from an anatomical MRI acquired either before or after birth [3]. Global shape descriptors such as the gyrification index can then be computed for each individual and compared across groups. Early disturbance in cortical development at birth in preterm babies can be quantified using this approach, and the extracted measures are predictors of infants' outcome at term equivalent age [5]. Global gyrification measures were also used in [4] to compare intra uterine maturation in fetuses and extra uterine maturation in age-matched preterm infants and showed that the folding dynamics is different.

II. FINER SCALE DESCRIPTORS

In order to get access to more local shape descriptors, we recently adapted a surface registration technique developed initially for adult brain to address the problem of defining spatio-temporal correspondences across cortical surfaces from a group of fetuses [6]. The analysis of local curvature showed that brain folding is a progressive process, and that large folds observed in adults do emerge from smaller indivisible entities. Based on this observation, we developed a technique to decompose the sulci into smaller entities corresponding to putative first sulcal folds, denoted as sulcal pits [7]. Using a closely related approach [8], Im and colleagues have shown that sulcal pits are valid descriptors of brain folding in fetuses, allowing to detect abnormal folding patterns [9]. On the other hand we have shown that sulcal pits are also suitable for characterizing postnatal brain maturation and abnormal folding in a group of children with Autism

spectrum disorder between 1.5 and 10 years old [10]. Finally, sulcal pits and corresponding shape descriptors can also be extracted from synthetic folded surfaces resulting from bio-mechanical modeling such as proposed in [12] and compared to real data. In the future, these predictive models will be decisive for enhancing our understanding of neural mechanisms resulting in abnormal cortical maturation in developmental brain disorders.

III. CONCLUSION

Quantitative local/regional descriptors of the geometry of the cortical surface have a great potential to highlight abnormalities in cortical folding pattern and to detect departure from typical developmental trajectories in the perinatal period, both before and after birth. They will be instrumental to uncover the biological mechanisms underlying the cortical maturation.

REFERENCES

- [1] J. Dubois, G. Dehaene-Lambertz, S. Kulikova, C. Poupon, P. S. Hüppi, and L. Hertz-Pannier, "The early development of brain white matter: a review of imaging studies in fetuses, newborns and infants.," *Neuroscience*, vol. 276, pp. 48–71, Sep. 2014.
- [2] F. Rousseau, C. Studholme, R. Jardri, and M. E. Thomason, "In Vivo Human Fetal Brain Analysis Using MR Imaging," in *Fetal Development*, 2016, pp. 407–427.
- [3] A. Makropoulos, S. J. Counsell, and D. Rueckert, "A review on automatic fetal and neonatal brain MRI segmentation," *Neuroimage*, no. June, pp. 1–18, Jun. 2017.
- [4] J. Lefèvre, et al., "Are Developmental Trajectories of Cortical Folding Comparable Between Cross-sectional Datasets of Fetuses and Preterm Newborns?," *Cereb. Cortex*, vol. 26, no. 7, pp. 3023–3035, Jul. 2016.
- [5] J. Dubois, et al., "Primary cortical folding in the human newborn: an early marker of later functional development," *Brain*, vol. 131, no. 8, pp. 2028–41, Aug. 2008.
- [6] G. Auzias, F. De Guio, A. Pepe, F. Rousseau, J.-F. Mangin, N. Girard, J. Lefèvre, and O. Coulon, "Model-driven parameterization of fetal cortical surfaces," in *IEEE 12th International Symposium on Biomedical Imaging (ISBI)*, 2015, pp. 1260–1263.
- [7] G. Auzias, L. Brun, C. Deruelle, and O. Coulon, "Deep sulcal landmarks: Algorithmic and conceptual improvements in the definition and extraction of sulcal pits," *Neuroimage*, vol. 111, pp. 12–25, May 2015.
- [8] K. Im, H. J. Jo, J.-F. Mangin, A. C. Evans, S. I. Kim, and J.-M. Lee, "Spatial distribution of deep sulcal landmarks and hemispherical asymmetry on the cortical surface.," *Cereb. Cortex*, vol. 20, no. 3, pp. 602–11, Mar. 2010.
- [9] K. Im, et al., "Quantitative Folding Pattern Analysis of Early Primary Sulci in Human Fetuses with Brain Abnormalities," *Am. J. Neuroradiol.*, vol. 38, no. 7, pp. 1449–1455, Jul. 2017.
- [10] L. Brun, G. Auzias, M. Viellard, N. Villeneuve, N. Girard, F. Poinso, D. Da Fonseca, and C. Deruelle, "Localized Misfolding Within Broca's Area as a Distinctive Feature of Autistic Disorder," *Biol. Psychiatry Cogn. Neurosci. Neuroimaging*, vol. 1, no. 2, pp. 160–168, Mar. 2016.
- [11] S. Takerkart, G. Auzias, L. Brun, and O. Coulon, "Structural graph-based morphometry: A multiscale searchlight framework based on sulcal pits," *Med. Image Anal.*, vol. 35, pp. 32–45, Jan. 2017.
- [12] T. Tallinen, J. Y. Chung, F. Rousseau, N. Girard, J. Lefèvre, and L. Mahadevan, "On the growth and form of cortical convolutions," *Nat. Phys.*, pp. 1–15, 2016.

G.A., S.T., C.D., O.C. and J.L. are with Aix Marseille Univ, CNRS, INT, Institut de Neurosciences de la Timone, Marseille, France.

F.R. is with IMT Atlantique, LaTIM U1101 INSERM, UBL, Brest, France.

N.G. is with CRMBM UMR 7339, Aix-Marseille Université, CNRS, France and APHM Timone, Service de Neuroradiologie Diagnostique et Interventionnelle, Marseille, France.

(Corresponding author e-mail: guillaume.auzias@univ-amu.fr)