



## Bulk evolution for MC@sHQ + EPOS & HADRONIZATION

EMMI RRTF - GSI July 2016

---

Marlene Nahrgang, Jörg Aichelin, Pol Bernard Gossiaux, Klaus Werner

July 18, 2016


Duke University & Subatech Nantes

**Previously: MC@sHQ + Kolb Heinz (RHIC & LHC)**



(« use at your own risks »)

# Organisers' suggestion



# Importance of realistic bulk evolution


... assumption: QGP is locally thermalized  $\Rightarrow$  heavy quarks scatter off thermal medium constituents.

... information about the local medium temperatures, (chemical potentials) and velocities can be obtained from fluid dynamical models.

... modern fluid dynamical simulations of HIC have been extremely successful in describing the bulk evolution, in particular identified  $p_T$  spectra and flow.

... any model with  $P(\Delta E)$  produces the generic  $p_T$  shape of  $R_{AA}$ , magnitude depends on details in the Eloss model IN COMBINATION with the bulk evolution model!

T. Renk, PRC85 (2012) arxiv:1112.2503





elliptic fireball evolution

VS.

(simple) hydrodynamics

P.B. Gossiaux et al., arXiv:1102.1114

# EPOS model




(pPb at  $\sqrt{s} = 5$  TeV, 0 – 20% centrality, left: subtract results for 60 – 80% centrality)

...is an event-by-event MC model for proton-proton, proton-nucleus and nucleos-nucleos collisions.

K. Werner et al., Phys.Rev. C89 (2014); Phys.Rev.Lett. 112 (2014); Phys.Rev. C85 (2012); Phys.Rev. C82 (2010); ...


# EPOS model - initial conditions

- Gribov-Regge multiple scattering approach with DGLAP parton ladders = pomerons
- saturation scale motivated by a CGC:  
 $Q_s \propto N_{\text{part}} \hat{s}^\lambda$   
 $N_{\text{part}}$  connected nucleons to pomeron,  $\hat{s}$  energy of pomeron,  $\lambda = 0.25$
- parton ladders are treated as classical, relativistic strings with longitudinal and transverse momentum
- high density of strings (= flux tubes) in AA collisions




# EPOS model - core-corona approach

- at  $\tau_0$  fluid (core) and escaping hadrons (incl. jet hadrons) (corona)
- criterion based on momentum and density of string segments




# EPOS model - core-corona approach

- at  $\tau_0$  fluid (core) and escaping hadrons (incl. jet hadrons) (corona)
- criterion based on momentum and density of string segments
- assuming fast equilibration for core  $\Rightarrow$  initial  $T^{\mu\nu}$  of the fluid
- fluctuating initial conditions  
 $\rightarrow$  one random event  
(30 – 50% central,  
 $\sqrt{s} = 2.76$  TeV, Pb+Pb)



# EPOS model - core-corona approach

- at  $\tau_0$  fluid (core) and escaping hadrons (incl. jet hadrons) (corona)
- criterion based on momentum and density of string segments
- assuming fast equilibration for core  $\Rightarrow$  initial  $T^{\mu\nu}$  of the fluid
- fluctuating initial conditions  
 $\rightarrow$  one random event  
(30 – 50% central,  
 $\sqrt{s} = 2.76$  TeV, Pb+Pb)



- core-corona separation:  $\rightarrow$  one random event

( $b = 5.7$  fm, pPb, positions of the projectile nucleon (yellow dot), the Pb surface (yellow line), the hit target nucleons (stars), Pomerons

(blue dots), as well as the core (red circles) and the corona (green circles) string segments.




# EPOS model - (viscous) fluid dynamical evolution

uses the 3 + 1 dimensional (viscous with  $\eta/s = 0.08$ ) fluid dynamical code by Iu. Karpenko, vHLLE [Comput.Phys.Commun. 185 \(2014\)](#)

initial mapping:

$$T^{\mu\nu}(x) = \sum_i \frac{\delta p_i^\mu \delta p_i^\nu}{\delta p_i^0} g(x - x_i)$$


- solves second-order viscous fluid dynamical equations
- provides space-time evolution of energy density and fluid velocities



parametrization of the lattice equation of state [WB collab., JHEP 1011 \(2010\)](#)

# EPOS model - freeze-out and hadronic afterburner


transverse momentum distributions ( $\pi$ ,  $K$ ,  $p$ ) for different centralities (EPOS2)



green: ALICE data    blue: without hadronic cascade    red: full calculations

- particlization on hypersurface of constant energy density/temperature  
 $T_H = 166$  MeV, Cooper-Fry procedure.  
(but space-time evolution of fluid dynamical fields is available until  
 $T \sim 130$  MeV)
- subsequent hadronic final state interactions via UrQMD.

# Particle production - corona vs. fluid



in pPb collisions



total

core - dominant at

low/intermediate  $p_T$


corona - dominant at high- $p_T$

# Flow in pPb collisions



mass splitting for  $v_2$  as in PbPb collisions!


# Flow in PbPb collisions



(EPOS2)


left:  $v_2$  with respect to opposite hemisphere sub-event plane

below:  $v_n$  from di-hadron correlations, semi-peripheral collisions, data from ALICE,  $p_{T,assoc} = 0.25 - 0.5$  GeV/c



# Epos model – Freeze out and hadronic afterburner


## Kolb-Heinz



Kolb Rapp (2003)

better agreement if  
initial flow ( $v_r = \tanh(0.02 r)$ )


## EPOS2.17V3




# Temperature profiles

A comparison of the temperature evolution in the middle of the fireball  
( $x = y = \eta = 0$ ):

RHIC,  $\sqrt{s} = 200$  GeV



LHC,  $\sqrt{s} = 2.76$  TeV




(EPOS2)


# Velocity profiles

A comparison of the radial velocities as a function of the radius in central collisions:

RHIC,  $\sqrt{s} = 200$  GeV





LHC,  $\sqrt{s} = 2.76$  TeV




(EPOS2 - includes preequilibrium flow)



# Velocity profiles (bis)



EPOS2 hotter and more explosive than Kolb-Heinz




# Hadronization

---

- Coalescence (Main ref.: PHYSICAL REVIEW C **79**, 044906 (2009) )
- Global hadronization algorithm

# 2 physical pictures for coalescence

Lin et Molnar (03) et de Fries,  
Müller, Nonaka and Bass (03)



2->1

C. B. Dover, U. W. Heinz, E.  
Schnedermann, and J.  
Zimanyi,  
Phys. Rev. C **44**, 1636 (1991).

Even for 2->1, several  
ways

$$N_{\Phi=D,B} = \int p_Q \cdot d\sigma_1 p_q \cdot d\sigma_2 \frac{d^3 p_Q}{(2\pi\hbar)^3 E_Q} \frac{d^3 p_q}{(2\pi\hbar)^3 E_q}$$

A delta distribution  $\longrightarrow \times f_Q(x_Q, p_Q) f_q(x_q, p_q) f_{\Phi}(x_Q, x_q; p_Q, p_q)$ .

# Dover vs Gossiaux (spatial)

**Dover**

Coall. Prob.: =  $D_{sp} \times D_p$

$$C(x_1, x_2; p_1, p_2) = \alpha_{1,2}^C \exp \left[ \frac{(x_1 - x_2)^2}{2R_c^2} \right] \times \exp \left[ \frac{(p_1 - p_2)^2 - (m_1 - m_2)^2}{2P_c^2} \right], \quad (2.14)$$

Particles far away can coalesce !!!

**Gaussian space (Gossiaux)**

$$f_\Phi(x_Q, x_q; p_Q, p_q) = \exp \left( -\frac{\|\vec{x}_{q,ref Q}\|^2}{2R_c^2} \right) \times \tilde{f}_\Phi(p_Q, p_q)$$

$$= \exp \left( \frac{\Delta x^2 - (\Delta x \cdot u_Q)^2}{2R_c^2} \right) \times \tilde{f}_\Phi(p_Q, p_q)$$

2.2.1 (Dover)

2.2.2 (Gossiaux)

$$N_\Phi(x_Q; p_Q) = g_q \int_{p_q \cdot \hat{d}\sigma(x_Q) > 0} \frac{d^3 p_q}{E_q} p_q \cdot \hat{d}\sigma(x_Q) e^{-\frac{u_{cell}(x_Q) \cdot p_q}{T}} \left( \frac{R_c}{\sqrt{2\pi\hbar}} \right)^3 \tilde{f}_\Phi(p_Q, p_q) \quad (29)$$

$$N_\Phi(x_Q; p_Q) = g_q \int_{p_q \cdot \hat{d}\sigma(x_Q) > 0} \frac{d^3 p_q}{E_q} \underbrace{p_q \cdot \hat{d}\sigma(x_Q)}_{u_Q \cdot \hat{d}\sigma(x_Q)} e^{-\frac{u_{cell}(x_Q) \cdot p_q}{T}} \left( \frac{R_c}{\sqrt{2\pi\hbar}} \right)^3 \tilde{f}_\Phi(p_Q, p_q) \quad (48)$$

The only difference

# Dover vs Gaussian (momentum)

## I. Gaussian p (2.3.1)

$$\tilde{f}_\Phi(\vec{p}_Q, \vec{p}_q) = \exp\left(-\frac{\|\vec{q}\|^2}{2\Lambda_{\text{cm}}^2}\right) = \exp\left(-\frac{\|\vec{p}_{q,\text{ref } Q}\|^2}{2m_q^2\alpha_g^2}\right)$$

(58)

$$\Lambda_{\text{cm}} = \frac{m_q m_Q}{m_q + m_Q} \alpha_g$$

(57)

Then (also using normalisation)

$$N_\Phi(x_Q; p_Q) = g_q \int_{p_q \cdot \hat{d}\sigma(x_Q) > 0} \frac{d^3 p_q}{E_q} p_q \cdot \hat{d}\sigma(x_Q) e^{-\frac{u_{\text{cell}}(x_Q) \cdot p_q}{T}} \left(\frac{R_c}{\sqrt{2\pi\hbar}}\right)^3 \tilde{f}_\Phi(p_Q, p_q) \quad (29/48)$$

$$\underbrace{\left(\frac{\sqrt{2\pi}(m_q + m_Q)}{m_Q m_q \alpha_g}\right)^3}_{c_g} \exp\left(-\frac{\|\vec{p}_{q,\text{ref } Q}\|^2}{2m_q^2\alpha_g^2}\right)$$

Dover (x) - Gaussian (p)

with  $\|\vec{p}_{q,\text{ref } Q}\|^2 = m_q^2 [(u_Q \cdot u_q)^2 - 1]$

$$N_\Phi(x_Q; p_Q) = \tilde{c}_g g_q \int_{u_q \cdot \hat{d}\sigma(x_Q) > 0} \frac{d^3 u_q}{u_0} u_q \cdot \hat{d}\sigma(x_Q) e^{-\left(\frac{m_q}{T} u_{\text{cell}}(x_Q) \cdot u_q + \frac{(u_Q \cdot u_q)^2 - 1}{2\alpha_g^2}\right)} \quad (60)$$

with  $\tilde{c}_g = \left(\frac{(m_q + m_Q)}{\sqrt{2\pi} m_Q \alpha_g}\right)^3$

Gossiaux (x) - Gaussian (p):  $u_q \cdot \hat{d}\sigma(x_Q) \rightarrow \frac{u_q \cdot \hat{d}\sigma(x_Q)}{u_Q \cdot \hat{d}\sigma(x_Q)}$

# Dover vs Gaussian (momentum)

## II Dover (2.3.2)

$$\tilde{f}_\Phi(p_Q, p_q) = \exp\left(-\left(\frac{m_Q m_q}{(m_q + m_Q)\Lambda_d}\right)^2 (u_q \cdot u_Q - 1)\right) \quad \text{with} \quad \Lambda_d = \frac{m_q m_Q}{m_q + m_Q} \alpha_d$$

(63)

Then (also using normalisation)  $\left(\frac{R_c}{\sqrt{2\pi\hbar}}\right)^3 = \left(\frac{m_Q + m_q}{m_Q m_q}\right)^3 \times \frac{1}{4\pi\alpha_d^2 K_2\left(\frac{1}{\alpha_d^2}\right) e^{\frac{1}{\alpha_d^2}}}$

$$N_\Phi(x_Q; p_Q) = g_q \int_{p_q \cdot \hat{d}\sigma(x_Q) > 0} \frac{d^3 p_q}{E_q} p_q \cdot \hat{d}\sigma(x_Q) e^{-\frac{u_{\text{cell}}(x_Q) \cdot p_q}{T}} \left(\frac{R_c}{\sqrt{2\pi\hbar}}\right)^3 \tilde{f}_\Phi(p_Q, p_q) \quad (29/48)$$

Dover (x) - Dover (p)

$$N_\Phi(x_Q; p_Q) = \tilde{c}_d g_q \int_{u_q \cdot \hat{d}\sigma(x_Q) > 0} \frac{d^3 u_q}{u_0} u_q \cdot \hat{d}\sigma(x_Q) e^{-\left(\frac{m_q}{T} u_{\text{cell}}(x_Q) + \frac{u_Q}{\alpha_d^2}\right) \cdot u_q} \quad (73)$$

with  $\tilde{c}_g = \left(\frac{m_Q + m_q}{m_Q}\right)^3 \times \frac{1}{4\pi\alpha_d^2 K_2\left(\frac{1}{\alpha_d^2}\right)}$

$$u_q \cdot \hat{d}\sigma(x_Q) \rightarrow \frac{u_q \cdot \hat{d}\sigma(x_Q)}{u_Q \cdot \hat{d}\sigma(x_Q)}$$

Gossiaux (x) - Dover(p) (our favorite)

4 combinations

$$N_{\Phi,D,G}(x_Q u_{\text{cell}} \cdot u_Q) = \tilde{c}_g g_q \int \frac{d^3 u_q}{u^0} u_q \cdot u_{\text{cell}} e^{-\left(\frac{m_q}{T} u_q \cdot u_{\text{cell}} + \frac{(u_Q \cdot u_q)^2 - 1}{2\alpha_g^2}\right)} \quad (75)$$

$$N_{\Phi,D,D}(x_Q; u_{\text{cell}} \cdot u_Q) = \tilde{c}_d g_q \int \frac{d^3 u_q}{u^0} u_q \cdot u_{\text{cell}} e^{-\left(\frac{m_q}{T} u_{\text{cell}} + \frac{u_Q}{\alpha_d^2}\right) \cdot u_q} \quad (76)$$

↑ ↑  
space momentum

$$N_{\Phi,Gx,G}(x_Q; u_{\text{cell}} \cdot u_Q) = \tilde{c}_g g_q \int \frac{d^3 u_q}{u^0} \frac{u_q \cdot u_{\text{cell}}}{u_Q \cdot u_{\text{cell}}} e^{-\left(\frac{m_q}{T} u_q \cdot u_{\text{cell}} + \frac{(u_Q \cdot u_q)^2 - 1}{2\alpha_g^2}\right)} \quad (77)$$

$$N_{\Phi,Gx,D}(x_Q; u_{\text{cell}} \cdot u_Q) = \tilde{c}_d g_q \int \frac{d^3 u_q}{u^0} \frac{u_q \cdot u_{\text{cell}}}{u_Q \cdot u_{\text{cell}}} e^{-\left(\frac{m_q}{T} u_{\text{cell}} + \frac{u_Q}{\alpha_d^2}\right) \cdot u_q} \quad (78)$$



One needs to fix the parameters ( $m_q$  and  $\Lambda_{\text{xxx}} / \alpha_{d/g}$ )

**Calibration:** section 2.4.5 (& 2.4.3)

**Assuming:**  $\mathbf{u}_{\text{cell}}$  oriented along  $d\sigma$  and general  $\mathbf{p}_Q$   
**And Gossiaux spatial**

Parameters tuned such that prob  $b \rightarrow B$  is = 1 through coalescence

**Dover p (set II)**



$m_q \approx 100 \text{ MeV}$  et  $\alpha_d = 0.876$  (set II<sub>b</sub> bis)

Optimal wrt hist. model



Dover spacial

$$N_{\Phi, D, D}(x_Q; p_Q) = \left( \frac{m_Q + m_q}{m_Q} \right)^3 \frac{m_q g_q}{\alpha_d^2 T K_2 \left( \frac{1}{\alpha_d^2} \right)} \times (u_{\text{cell}} + \mu u_Q) \cdot \hat{d}\sigma \times \frac{K_2(D)}{D^2}$$

with  $D = \frac{m_q}{T} \times \sqrt{1 + 2\mu u_{\text{cell}} \cdot u_Q + \mu^2}$  et  $\mu = \frac{T}{m_q \alpha_d^2}$

Gossiaux spatial

$$N_{\Phi, Gx, D}(x_Q; p_Q) = \left( \frac{m_Q + m_q}{m_Q} \right)^3 \frac{m_q g_q}{\alpha_d^2 T K_2 \left( \frac{1}{\alpha_d^2} \right)} \times \left( \mu + \frac{u_{\text{cell}} \cdot \hat{d}\sigma}{u_Q \cdot \hat{d}\sigma} \right) \times \frac{K_2(D)}{D^2}$$

After normalization

Depends on 3 invariant quantities

$$\text{prob}_{\Phi}(x_Q; p_Q) = \max \left( f f \times \text{prob}_{\Phi}(x_Q; p_Q, \hat{d}\sigma = u_{\text{cell}}), 1 \right)$$

$$\text{prob}_{\Phi}(x_Q; p_Q, \hat{d}\sigma = u_{\text{cell}}) = \left( \mu + \frac{1}{u_Q \cdot u_{\text{cell}}} \right) \left( \frac{m_Q + m_q}{m_Q} \right)^3 \frac{m_q g_q}{T} \frac{K_2(D)/D^2}{K_2 \left( \frac{1}{\alpha_d^2} \right) / \frac{1}{\alpha_d^2}}$$

Fig 2 of PRC79

$$f f = \frac{\mu + \frac{u_{\text{cell}} \cdot \hat{d}\sigma}{u_Q \cdot \hat{d}\sigma}}{\mu + \frac{1}{u_Q \cdot u_{\text{cell}}}}$$

Flux factor

# Caviats


- 1)  $m_q$  probably too small
- 2) Coalescence -> « average » D meson (chemistry not taken into account)

# Hadronization algorithm


- 1) For pp: use  $p_T$ -frag ( $y_D=y_c$  and  $\mathbf{p}_{TD}=z \mathbf{p}_{Tc}$ )
- 2) For pA or AA:
  - 1) Perform criteria in the fluid cell rest frame
  - 2) When fragmentation is chosen perform the same fragmentation as in pp, staying in the lab frame

This allows to preserve the rapidity invariance as well as  $Q$  (ratio) =1 if no Eloss (and no shadowing)...(NOT SO TRIVIAL !!!) Then looking for the coal + frag case:

$dN_D/dp_T$



$Q_D(p_T)$





While boost invariance is preserved as well for boost invariant QGP (checked but not shown) 31

**some effects on HQ evolution**

---

# Basic observables (RHIC)



## Kolb Heinz hydro



Large effects from coalescence (especially in the  $R_{AA}$ )... to be studied in a more systematic way

# Basic observables (LHC 2.76 TeV)

EPOS2




Large effects from coalescence for both observables

# Comparison between EPOS2 and EPOS3

EPOS2: ideal fluid dynamics with initial smearing to flux tube size to mimick viscosity

EPOS2+shadowing

EPOS3: viscous fluid dynamics ( $\eta/s = 0.08$ ) and individual parton saturation scales as described earlier



EPOS3 calculations by V. Ozvenchuk

⇒ visible difference even in two similar approaches that both describe light hadron data VERY WELL... problematic or could be an opportunity to probe more details?!


# Average vs. fluctuating IC

- consider medium with local fluctuations  $\varepsilon(x) = \varepsilon_0 + \delta\varepsilon(x)$  around an averaged energy density  $\langle\varepsilon(x)\rangle = \varepsilon_0$
- local energy loss:  $dE/dx \propto T^\beta$  with  $\beta \approx 1$  for collisional energy loss with running  $\alpha_s$  to  $\beta \approx 2$  for radiative LPM

- $\varepsilon \propto T^n$ , with  $n \approx 4$  from EoS

- $\frac{dE}{dx} \propto \varepsilon^\delta$  with  $\delta = \frac{\beta}{n} < 1$

$$\left\langle \frac{dE}{dx} \right\rangle = \frac{dE(\varepsilon_0)}{dx} \times \left[ 1 - \frac{\delta(1-\delta)}{2} \frac{\langle \delta\varepsilon^2 \rangle}{\varepsilon_0^2} + \dots \right].$$





⇒ Less energy loss for event-by-event studies compared to average medium evolution!



# Fluctuating IC $\rightarrow$ finite triangular flow

Fluctuating initial conditions enables us to study higher order flow harmonics for heavy quarks:






$\Rightarrow$  finite triangular flow for D mesons predicted! First STAR analysis under way.

M. Nahrgang, J. Aichelin, S. Bass, P. B. Gossiaux, K. Werner, Phys.Rev. C91 (2015); J.Phys.Conf.Ser. 668 (2016)

# Interpretation of the EoS

scattering partner of HQ are thermal quasiparticles in the medium  $\Rightarrow$  nature of these quasiparticles matter!

typically, one uses either massless quasiparticles or with thermal masses from pQCD, which DOES NOT reproduce the lattice EoS!



# Conclusions

The use of a realistic model for the bulk evolution is indispensable for the study of traditional ( $R_{AA}$  and  $v_2$ ) and new ( $v_n$  and  $\Delta\phi$  correlations) HQ observables!

It can also serve as an opportunity to investigate more details of the QGP: initial conditions, nature of quasiparticles, etc.

**Back up**



---

# Some EPOS2+MC@sHQ results at LHC

K values fixed at  $p_T=10$  GeV/c, x2 if reduction of dof according to EOS134 !

Still close to unity if rad + col

considered





Large push from the radial flow; discrepancy unlikely to be explained by shadowing alone.

Concerns: Need to revisit the model for small  $p$  ? (Bad) consequences for  $v_2$  ?

**Main message: the models of HF energy loss and the background medium (including its microscopic content) are bound together**

# Some EPOS2+MC@sHQ results at RHIC



Larger radial flow in EPOS

Both « cocktails » (HF energy loss + background + K factor) provide a fair agreement with the data

Data at larger  $p_T$  would help a lot !

**Main message: the models of HF energy loss and the background medium (including its microscopic content) are bound together**