





Dosimetric environment and first results of the studies on the physicochemical and biological mechanisms of FLASH hadrontherapy at the ARRONAX cyclotron YARRONAX



de l'Ouest

unicancer

PAYS DE LA LOIRE



Manon Evin, Charbel Koumeir, Quentin Mouchard, Guillaume Blain, Emeline Craff, Gregory Delpon, Vincent Fiegel, Giovanna Rosa Fois, Youssef Ghannam, Ferid Haddad, Lydia Maigne, Vincent Potiron, Gaëlle Saade, Noël Servagent, Stéphane Supiot, Sarra Terfas, Johan Vandenborre, Vincent Métivier, Sophie Chiavassa



FLASHMOD project



Multidisciplinary team around the ARRONAX cyclotron

Accélérateur pour la Recherche en Radiochimie et Oncologie à Nantes-Atlantique



Extracted	Maximal	Range in	LET at plateau
particles	energy (MeV)	water (cm)	entrance $(keV/\mu m)$
H^{+}	70	4,08	1
He^{2+}	70	0, 34	11



Up to several hundred of **kGy/s**



• Poirier F et al., « The injection and chopper-based system at ARRONAX C70XP cyclotron », cyc19, sept. 2019, doi:10.18429/jacow-cyclotrons2019-tup006

Dosimetric environment

Applications

UHDR and CONV beam monitoring

and the state of the

Quentin Mouchard



• Mouchard Q., et al.. Upgrade of the flash beam monitoring system at ARRONAX cyclotron. (FRPT 2022). (hal-03885576)

In vivo dosimetry with radiochromic films

LET (Linear Energy Transfert) dependency







Gate simulation

→ Determination of the correction factor **f(LET)**

- 12 %

Context	Dosimetric environment		Applications		Perspectives	
Radiobiology:		✓ Biolog✓ Rapid	gically complex growth			
> Dose rate effect on	healthy tissues	Irradiation	\rightarrow	 ✓ High s Study of t the FLASH 	tatistical power he irradiation para l effect	ameters triggering
Protocol	Assembling the Hypoxia Bragg plateau	FE DE CE BE AT	Culture (28°C)	4 days A al	nesthesiea (cold) nd fixation	Acquisition of images and analysis
Results Length Curvature Pericardial oedema 	Control (unirradiated)	7500 Gy/s	D,25 Gy/s	✓ P Saadé	Protective effect with H ⁺ et Mechanis Biological	ct of FLASH He ²⁺ ms ? Radiochemical

• Saadé et al., « Ultrahigh-Dose-Rate Proton Irradiation Elicits Reduced Toxicity in Zebrafish Embryos, Advances in Radiation Oncology, 2022. DOI: 10.1016/j.adro.2022.101124

• Ghannam Y. et al., « First evidence of in vivo effect of FLASH radiotherapy with helium ions in zebrafish embryos », 2023. <u>https://doi.org/10.1016/j.radonc.2023.109820</u>.



Dose rate (Gy s⁻¹)

30

- 28 %

-27%

OC-1 films

Radiochemistry: experimental Irradiation Protocol $ROS \rightarrow$ indirect effects of radiations Reactive oxygen species produced by the radiolysis of **PEEK cells** Simulation (GATE) Normalized dose water Measurement Ultrapure water Atmospheres : 1. Air $[H_2O_2]$ when the dose rate increases / 2. N₂O 3. Argon Mechanism ? 10 20 Depth in water (mm) Protons in plateau Main production by Results $H0^{\bullet} + H0^{\bullet} \rightarrow H_2 0_2$ H_2O_2 yield at 80 Gy at variable dose rates G_{TS}(H₂O₂) (×10⁻⁷ mol J⁻¹) 0.0 0.1 2.1 7.7 Recombination process of HO°? $\rm H0^{\bullet}$ + $e_{aq}^{-} \rightarrow \rm H0^{-}$ Privileged at UHDR? CONV UHDD Under N_2O : no e_{aq} , no O_2 in solution UNDER N2O UNDER AIR UNDER ARGON 0.0 Under Argon: no O₂ in solution 10 100 1000 Ω 1

1. Key role of e_{aq}^- at the radiolysis step

Spectro-photometry

UV-Visible

2. O_2 depletion has a minor impact

- Blain, et al.. « Proton Irradiations at Ultra-High Dose Rate vs. Conventional Dose Rate: Strong Impact on Hydrogen Peroxide Yield. » Radiation Research, 2022, 198, pp.318-324. (10.1667/RADE-22-00021.1)
- Fiegel et al., « Investigation of the radiolytic ROS production mechanism under proton FLASH conditions: evidence of the key role of the aqueous electron. », submitted.



1

0,5

1.E-03

1.E-01

1,E+01

1,E+03

1,E+05

Time in ns

1.E+07

1,E+09

- dosimetry)
- Work in progress to model UHDR

· Giovanna Rosa Fois, et al.. « Microdosimetry and radiolytic species production in UHDR proton beam using GATE and Geant4-DNA. » (2022) Napoli, Italy. (hal-03886937)



Future application: mouse irradiations



Study of preclinical set-up and instrumentation



Methodology for:

• Optimal field size with fixed collimators

Optimal homogeneous field size (cohort) = Organ size (cohort) + Margins Targeting + Margins Respiratory

• Internal organs targeting without online imaging



- *In vivo* dosimetry with radiochromic OC-1 films
- Impact of the respiratory cycle on the dose distribution on FLASH-RT

In vivo dosimetry with radiochromic films

Mouse whole brain irradiation: simulation from XRAD225Cx µCBCT image and targeting without image guidance at ARRONAX

Simulation

Measure DD

Dose





Distance

1D illustration of the gamma index metric

Simulated



Local gamma index 2D (DD 1% - DTA 1 mm): pass rate 98,2 %



9

Impact of the respiratory cycle in FLASH-RT

FLASH-RT irradiation time ~ ms

Dosimetric impact ?

MOBY (Mouse Whole Body 4D Digital Fantom) V.2

Delta AP



Full expiration and inspiration (density, composition and volume)

 Heyden, Brent van der, et al.. « The Influence of Respiratory Motion on Dose Delivery in a Mouse Lung Tumour Irradiation Using the 4D MOBY Phantom ». *The British Journal of Radiology* 90, n° 1069 (janvier 2017): 20160419. <u>https://doi.org/10.1259/bjr.20160419</u>.

Fluoroscopic mode XRAD225Cx (8 images/s)



Maximal amplitudes over 10 cycles of an anaesthetized mouse



Impact of the respiratory cycle in FLASH-RT

Lateral irradiation



Full expiration (Deflate) (d=0,45 g/cm³, volume 0,56 cm³)



Lateral (left side) irradiation

Right





Differences in homogeneity and mean dose:

- < 1 % for entrance lung
- [1-3] % for exit lung



Vertical irradiation







Conclusion and perspectives



Demonstration of the effect of the dose rate on zebrafish embryos and in vitro
 Advances in physico-chemical mechanisms and development of a simulation code

- Zebrafish embryos: analysis of gene activation
- ---> Radiochemistry: radiolysis in biological media
- ---> Irradiation of small animals (tumors and healthy tissue)
- → Design of a ridge filter for irradiation in the Spread Out Bragg Peak (SOPB)



